Multi-Scale Model-Driven Sampling with Autonomous Systems At A National Littoral Laboratory: Turbulence Characterization from an AUV

Edward R. Levine Naval Undersea Warfare Center, Division Newport Code 8211 Newport, RI 02841-1708

phone: (401) 832-4772 fax: (401) 832-2146 email: levineer@tech.npt.nuwc.navy,mil Award #: N000149WX20272, N000149WX30337 http://marine.rutgers.edu/MAS

LONG-TERM GOAL

The long term goal of this research is to utilize turbulence measurements obtained from small Autonomous Underwater Vehicle (AUV) based sensors as the subgrid characterization tool in coastal ocean observation/prediction networks.

OBJECTIVES

The objective of this research is to use AUV-based turbulence measurements to quantify mixing in shallow water physical process studies (upwelling regions, fronts. boundary layers) within the context of the LEO-15 based National Ocean Partnership Program (NOPP) coupled ocean observation/modeling system This includes estimating mixing levels, identifying regions of enhanced mixing, determining the horizontal spatial scale of mixing events, defining the role of boundary layers, and parameterizing results for coastal predictive model testing studies of subgrid scale processes.

APPROACH

The initial stage of the approach is to integrate an optimum turbulence sensor suite into a small, logistically simple, AUV, with input from the ocean turbulence and modeling communities. The next stage is to establish this small AUV as a viable platform for coastal turbulence research. Towards this end, horizontals profiles of estimated dissipation rate, temperature microstructure, 3-dimensional small scale velocity, finescale vertical shear of horizontal current, and stratification are obtained in the coastal environment.

Subsequently, mixing is studied in the context of the multi-scale measurements surrounding the NOPP LEO-15 node site in the Mid-Atlantic Bight, during the summer upwelling season. Sampling is done adaptively using input from the Rutgers University continental shelf model SCRUM (Song and Haidvogel, 1994). The sensor suite enables estimates of eddy diffusivity profile (Gargett and Moum (1995), eddy viscosity profile (using the truncated TKE equation), Richardson numbers, and fluxes [using the correlation technique]. These data enable us to evaluate turbulence closure schemes associated with subgrid mixing processes as parameterized by 3 different surface boundary layer submodels in SCRUM, the coastal circulation model

Public reporting burden for the col maintaining the data needed, and c including suggestions for reducing VA 22202-4302. Respondents shot does not display a currently valid C	ompleting and reviewing the collecthis burden, to Washington Headquild be aware that notwithstanding a	tion of information. Send comment parters Services, Directorate for Inf	s regarding this burden estimate formation Operations and Reports	or any other aspect of the s, 1215 Jefferson Davis	his collection of information, Highway, Suite 1204, Arlington	
1. REPORT DATE 30 SEP 1999 2. REPORT TYPE			3. DATES COVERED 00-00-1999 to 00-00-1999			
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER		
Multi-Scale Model-Driven Sampling with Autonomous Systems At A National Littoral Laboratory: Turbulence Characterization from an				5b. GRANT NUMBER		
AUV				5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)				5d. PROJECT NUMBER		
				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Undersea Warfare Center, Division Newport,Code 8211,Newport,RI,02841				8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)		
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAIL Approved for publ		ion unlimited				
13. SUPPLEMENTARY NO	TES					
14. ABSTRACT						
15. SUBJECT TERMS						
16. SECURITY CLASSIFIC		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON		
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	5		

Report Documentation Page

Form Approved OMB No. 0704-0188

WORK COMPLETED

The REMUS AUV has been instumented with a turbulence sensor package, which has been cantilevered off the bow or integrated into the hull. (Levine and Lueck, 1999) (Fig. 1). Sensors include two shear probes, an ultra-fast thermistor, an upward and downward looking ADCP, two CTDs, and an ADV-O.




Fig 1. The REMUS AUV instrumented with turbulence sensors

Utilizing this system, scientific studies of the Ocean Boundary Layer (OBL) during an upwelling event were conducted near the Rutgers University LEO-15 site on the inner continental shelf off New Jersey during July 1998. In the field experiment, high quality data were obtained from all sensors, and a turbulence characterization in various components of the upwelling system has been obtained.

RESULTS

Model-based adaptive sampling, updated with synoptic information from the other platforms, was used to deploy the AUV was along trajectories through components of the Leo-15 upwelling region, including the upwelling gyre and offshore jet. Model predictions include assimilated data from the wide variety of sampling platforms which characterize larger scale structures on the continental shelf.

Results indicate that the modified REMUS AUV was a viable platform for turbulence data acquisition in the coastal ocean. Analysis of the shear probe data indicate that after platform noise removal, using the techniques of Levine and Lueck (1999), the shear autospectra agree with the Nasmyth "universal spectrum" (Oakey, 1982) out to wavenumbers close to the physical size of the sensing tip of the probes.

For the case of fully developed upwelling , an example of the upwelling gyre center and offshore jet turbulence characterization estimation is shown in Fig 2. For a 5 m depth transit, time series of mixing parameter estimates primarily show dissipation rates of 10^{-7} to 10^{-6} W kg $^{-1}$, eddy diffusivities of 10^{-5} to 10^{-4} m 2 s $^{-1}$, eddy viscosities of 10^{-4} to 10^{-3} m 2 s $^{-1}$, and Richardson numbers of 10^0 to 10^1 .




Fig 2. Mixing parameters from LEO-15 upwelling region, July 1998

IMPACT/APPLICATION

The AUV-based turbulence measurements provide a unique horizontal profiling view of the variability of the mixing environment that cannot be obtained by more conventionally sampling measurements, and this approach can be further exploited in yo-yoed horizontal sections. These techniques are invaluable in upwelling process studies in which competing turbulence closure model alternatives are

testing in SCRUM to parameterize subgrid processes. Competing OBL alternatives include those of Mellor-Yamada (1974) Large et al. (1994), and Price et al. (1986). Features such as the evolution of the upwelling front can be tested.

TRANSITIONS

Our AUV sensor technologies, hardware and software, are being considered for inclusion as tactical oceanography payloads for the Manta UUV Initiative

RELATED PROJECTS

Follow-on studies with the Rutgers group were conducted in 1999 under separate NOPP support. Also, my AUV-based turbulence measurement system is being utilized in NOPP studies with the Harvard led LOOPS project, for which measurements were also made in Cape Cod Bay during September 1998. The system is also being utilized in NOPP FRONT studies on the New England continental shelf during 1999-2001.

REFERENCES

- A. E. Gargett and J. N. Moum. 1995. Mixing effects in tidal fronts: results from direct and indirect measurements of density flux. J. Phys. Ocean., 25, 2583-2608.
- W. G., Large, J. C. McWilliams, and S. C. Doney. 1994: Oceanic vertical mixing: a review and model with a non-local boundary parameterization. Rev. Geophys., 32, 363-403.
- Levine, E. R., R. G. Lueck, 1999. Turbulence measurements with an autonomous underwater vehicle. Journal of Atmospheric and Oceanic Technology, Special Issue on Ocean Turbulence Measurement, 16, 11, part 1, 1533-1544.
- G. L. Mellor and T. Yamada. 1974. A hierarchy of turbulence closure models for planetary boundary layers. J. Atmos. Sci., 31, 1791-1806.
- N. S. Oakey. 1982. Determination of the rate of dissipation of turbulent energy from simultaneous temperature and velocity shear measurements. J. Phys. Ocean., 12, 256-271.
- J. F Price., R. A. Weller, and R. Pinkel. 1986. Diurnal cycling: Observations and models of the upper ocean response to diurnal heating, cooling, and wind mixing. J. Geophys. Res., 91, C7, 8411-8427.
- Y. Song and D. Haidvogel. 1994. A semi-implicit ocean circulation model using a generalized topography following coordinate system. J. Comput. Phys., 115, 228-244.

PUBLICATIONS

Glenn, S. M., D. B. Haidvogel, O. M. E. Scofield, C J. von Alt, and <u>E. R. Levine</u>, 1998: Coastal Predictive Skill Experiments. Sea Tech., April 1998, 66-72.

- Levine, E. R., R. G. Lueck, 1998: A small AUV-based turbulence measurement system for NOPP combined coastal observation/prediction networks. EOS, Trans. Am. Geophys. Un., 1998 Ocean Sciences meeting, San Diego, Ca, Feb, 1998.
- Levine, E. R., R. G. Lueck, 1999: Turbulence measurements with an autonomous underwater vehicle. Journal of Atmospheric and Oceanic Technology, Special Issue on Ocean Turbulence Measurement, 16, 11, part 1, 1533-1544.
- Levine, E. R., R. G. Lueck, R. R. Shell, and P. Licis, 1999: Ocean turbulence estimates in the Coastal Ocean Modeling and Observation Program (COMOP) studies at LEO-15. ASLO 1999 Aquatic Scienes Meeting, Santa Fe. NM, Feb 1999.
- Levine, E. R., R. G. Lueck, R. R. Shell, and P. Licis, 1999: Coastal turbulence estimates and physical process studies utilizing a small AUV. Invited paper at 1999 Spring AGU meeting, Boston, MA, June 1999. abstract in. supple. to April 17 EOS, Trans. Am. Geophys. Un., S192.
- Levine, E. R., R. G. Lueck, R. R. Shell, and P. Licis, 1999: Coastal turbulence estimates and physical process studies utilizing a small AUV, Proceedings, Eleventh International Symposium on Unmanned Untethered Vehicle Technology (UUST99), Durham, NH. 94-102, Sept. 1999.
- Levine, E. R., R. G. Lueck, R. R. Shell, and P. Licis, 2000: Coastal Turbulence Estimates in Ocean Modeling and Observation Studies Near LEO-15, accepted for 2000 Ocean Sciences Meeting, Special Session on Coastal Ocean Dynamics and Prediction, Jan 2000, San Antonio. TX.